



Fast Regeneration
Restored Motion

Table of Contents

Burns

Hand scar revision after burn, <i>L. Kamolz, MD, Graz, Austria</i>	4
Pediatric scald burn, <i>J. Lee, MD, Seoul, S. Korea</i>	6
Massive scar contracture on flank, <i>J. Grünert, MD, St. Gallen, Switzerland</i>	8
Revision surgery of burn scar at the lower limb, <i>A. Modarressi, MD, Geneva, Switzerland</i>	10
Flame burn on chest and neck, <i>E. Dantzer, Toulon, France</i>	12

Trauma

Degloving of the forearm in a work accident, <i>M. Kerl, MD, Graz, Austria, Epigard by Biovision GmbH</i>	14
Degloving injury at lower leg, <i>M. Öhlbauer, MD, Murnau, Germany</i>	16
Crush trauma by a school bus, <i>M. Öhlbauer, B. Wallner and M. Miltz, MD, Murnau, Germany</i>	18
Avulsion injury at the foot with exposed Achilles tendon, <i>U. Graf, MD, Zurich, Switzerland</i>	20
Infected dog bite at the dorsum of the hand, <i>U. Hug, MD, Luzern, Switzerland</i>	22
Deglovement instep and plantar regions after motor-bike injury, <i>M. Öhlbauer, B. Wallner and M. Miltz, MD, Murnau, Germany</i>	24
Circular saw injury, <i>L. Mathys, MD, Bern, Switzerland</i>	26
Trauma by car accident, <i>D. F. Kalbermatten, MD, Basel, Switzerland</i>	28

Chronic Wounds

Chronic venous ulcer with 3.5 years follow-up, <i>F. Tostes, MD, Rio Grande do Sul, Brazil; V.A.C. therapy and GranuFoam by Kinetic Concept, Inc (KCI)</i>	30
Diabetic foot syndrome with gangrene, <i>D. Luedi, MD, Langenthal, Switzerland</i>	32
Chronic diabetic ulcer at clubfoot, <i>H. Mössner, MD, Salzburg, Austria; Medihoney by Derma Sciences, Inc; Hyiodine by Contipro Group s.r.o.</i>	34
Chronic mixed ulcer in diabetic patient, <i>M. Öhlbauer, MD, Murnau, Germany</i>	36
Large pressure ulcer after an ischaemic stroke, <i>M. Öhlbauer, MD, Murnau, Germany</i>	38

Cancer Excision

Large squamous cell carcinoma, <i>D. Luedi, MD, Langenthal, Switzerland</i>	40
Basal cell carcinoma excision in face, <i>N. Lilgenau, MD, Vienna, Austria</i>	42
Carcinoma resection in 97-year-old patient, <i>V. Wedler, MD, Frauenfeld, Switzerland</i>	44
Melanoma on the nasal ridge, <i>M. Öhlbauer, MD, Murnau, Germany</i>	46

Adhesion Barrier

Adherent scar after extravasation of cytotoxic drugs, <i>Keck et al. 2008, Handchir Mikrochir Plast Chir; ©Thieme</i>	48
Treatment of adherent foot tendon, <i>U. Graf, MD, Zurich, Switzerland</i>	50
Adherent scar at knee, <i>W. Meyer, MD, Fort Malakoff Clinic, Mainz, Germany</i>	52
Exposed tendon of the hand, <i>J. Federbusch, MD, Rio Grande do Sul, Brazil</i>	54

Donor Sites

Donor Site after Phalloplasty with the Radial Forearm Flap, <i>W. Watfa, MD, Lausanne, Switzerland</i>	56
--	----

Hand scar revision after burn

Type of Wound	Contractile scar revision after burn Injury*
Etiology	Hypertrophic and contractile scar after 3 rd degree flame burn of the right hand. Pronounced hyperextension of the metacarpophalangeal (MCP) joint and complete stiffness of the wrist.
Patient	Young female

Pre-operative: Hyperextension of the MCP joint – function is considerably limited (Fig.1). Fist closure is not possible (Fig.2). Intra-operative: Complete excision of the scar tissue (Fig.3). Intra-operative: The scar tissue – laid back onto the wound bed for the photo – reveals the dimension of the scar contraction, the skin deficit and the substantial functional limitation (Fig.4). Long-term follow-up: Normal function of the hand without any limitations (Fig.5–7).



Pediatric scald burn

Type of Wound	Scald burn on the foot*
Etiology	Burn wound
Patient	2-year-old child

A 2-year-old child presented with a scald burn on the right foot (Fig. 1). At the subsequent surgery the wound was carefully debrided (Fig. 2). Due to the higher risk of contracture because of the young age of the patient, it was decided to use MatriDerm® in an attempt to prevent scar tissue formation and contracture.

MatriDerm® was applied to the wound bed and trimmed appropriately (Fig. 3). A split-thickness skin graft (STSG) was placed on top of MatriDerm® in a One-Step Procedure (Fig. 4). The STSG was secured using staples.

At 2 days follow-up the wound showed good signs of healing and good take rate of the graft (Fig. 5). At 6 months the wound was completely healed without scarring (Fig. 6). At 1 year follow-up the skin demonstrated excellent pliability and good aesthetic result (Fig. 7). At 4 year follow-up the grafted area showed same skin color as the surrounding skin (Fig. 8). The regenerated skin is growing at the same rate as the surrounding tissue.

This case demonstrates that MatriDerm® can be used successfully in pediatric patients with excellent aesthetic and functional outcomes.



Massive scar contracture on flank

Type of Wound	Burn scar revision*
Etiology	Scar contracture of burn wound on flank
Patient	56-year-old male

The patient presented with a large scar from a burn wound on the flank 20 years ago (Fig. 1). Reason for his request of scar revision was the instability and aesthetic appearance of the scar and skin insensitivity. Comorbidities included a history of peripheral artery occlusive disease and arterial hypertonia.

The scar plate was excised completely and the contractures were released (Fig. 2). The resulting defect was covered with multiple dry sheets of MatriDerm® (Fig. 3), which were rehydrated in situ (Fig. 4).

A split-thickness skin graft (STSG) was positioned over MatriDerm® in a One-Step Procedure. Negative Pressure Wound Therapy (NPWT) was initiated to fixate the graft. After 5 days, the first dressing change was

performed and NPWT was discontinued. The wound showed a good take rate of the STSG, with just minimal dehiscence (Fig. 5).

Three weeks post-operatively, the graft was almost completely healed, demonstrating markedly reduced tension (Fig. 6). One month p.o., the wound was 90% tension free, with just a few small wounds (Fig. 7).

Six months p.o., there was no loss of functionality, with just a small scar cord in the axilla, that does not trouble the patient. Despite the still lowered mamilla, the patient is satisfied with the current result (Fig. 8).



Revision surgery of burn scar at the lower limb

Type of Wound	Burn sequela – unstable scar and functional impairment due to contractures*
Etiology	Flame burn in childhood (at age 8)
Patient	27-year-old female

Aspect of the wound before operation (Fig.1,2). Pre-operative marking of the area to be excised (Fig.3). Aspect of the wound after scar resection with release of healthy tissue. In order to show the extend of contracture release the scar tissue was put back onto the wound bed for this photo (Fig.4). Application of

MatriDerm® (Fig.5), covered with fenestrated split-thickness skin graft (Fig.6). Negative pressure wound therapy as dressing (Fig.7). Result after 2 weeks (Fig.8). Result after 3 months (Fig.9). Result 1 year later (Fig.10,11). Pleasing functional and aesthetic outcome with full pain release. Complete and stable healing of the wound.



Flame burn on chest and neck

Type of Wound	Flame burn on chest and neck*
Etiology	Burn wound
Patient	42-year-old male



A 42-year old male presented with a flame burn on chest and neck (60% TBSA) (Fig. 1). This is a challenging region due to high incidents of contractions, so high elasticity is very important.

The wound was carefully debrided to ensure a vital wound bed. MatriDerm® (1 mm) was applied to the wound bed (Fig. 2).

A split-thickness skin graft (STSG) was placed on top of MatriDerm® in a One-Step Procedure (Fig. 3).

Two and five years post-operatively, the grafted area showed very good elasticity and skin colour.

The good functional outcome was demonstrated by the unlimited extension of the neck (Fig. 4–7). This case demonstrates that MatriDerm® can be used successfully in demanding region like the neck with excellent aesthetic and functional outcome.



Degloving of the forearm in a work accident

Type of Wound	Avulsion injury*
Etiology	Avulsion with degloving of the upper extremity with release of compartments
Patient	20-year-old male

The patient suffered from a work accident where his arm was crushed into an industrial roller, which resulted in multiple bone fractures and degloving of his right arm (Fig. 1).
Two days after the accident a fasciotomy was performed to release the compartment syndrome of the forearm and hand (Fig. 2). The defect was covered with Epigard to clean and condition the wound bed (Fig. 3).
Twelve days after the accident the debrided wound was covered with dry sheets of MatriDerm®, which were cut into shape to fit within the wound bed. MatriDerm® was rehydrated with sterile physiological

saline (Fig. 4), immediately followed by placement of a meshed split-thickness skin graft on top of MatriDerm® in a One-Step Procedure (Fig. 5). Negative pressure wound therapy was initiated post-surgery.
At the follow-up visit three weeks later, stable wound closure with complete integration of the skin graft was observed (Fig. 6, 7). The patient had good initial mobility of his elbow and wrist. Five months later the patient was back at work.



* Courtesy of M. Kerl, MD, Graz, Austria
Epigard by Biovision GmbH


Degloving injury at lower leg

Type of Wound	Significant soft tissue loss of the left lower leg and foot with visible tendons and periosteal structures of the medial ankle*
Etiology	Degloving injury
Patient	71-year-old male

This 71-year-old patient suffered a degloving injury of the left lower leg and foot (Fig.1). After operative debridement, there was significant soft tissue loss with visible tendon and periosteal structures of the medial ankle (Fig.2). Angiography showed that the lower leg and foot were only being nourished by an arteriosclerotic tibialis posterior artery.

Defect coverage was performed using 1 mm MatriDerm® (Fig. 3) in a One-Step Procedure with unmeshed split skin grafts in combination with one week of negative pressure wound therapy to fix the grafts (Fig. 4).

With this procedure the leg could be saved and showed a stable wound closure – one month p. o. (Fig.5). Two years after the accident the patient was able to wear normal shoes and clinical gait analysis demonstrated a perfect functional outcome (Fig.6). The patient refused to have any further procedure done to improve the aesthetic outcome.




Crush trauma by a school bus

Type of Wound	Deep soft tissue defect with involvement of the fascia, right lower leg*
Etiology	Crush trauma by a school bus
Patient	7-year-old boy

The patient was transferred to a special trauma clinic on day 5 post injury. Two-thirds of the proximal lower leg showed necrotic tissue (Fig.1). Parts of the lower leg fascia were destroyed. After debridement vital paratenon at the tibia was exposed (Fig.2). The tissue defect was covered with 1mm MatriDerm® and unmeshed split-thickness skin graft in a One-Step Procedure (Fig.3,4). Fixation was performed by negative pressure wound therapy for 1 week. Thereafter conventional dressing (fatty gauze/bulky gauze/tight bandaging)

was performed and the wound healed quickly without any complications. Aftercare with compression garments. Good functional and aesthetic result was achieved. Fig.5: 18 months after injury. The transplanted skin was pliable and soft so that the patient was able to return to normal life. 7 years after the surgery the aesthetic outcome has further improved (Fig.6). No scar contracture release surgeries have ever been necessary because the new skin has grown naturally along with the boy's legs.



Avulsion injury at the foot with exposed Achilles tendon

Type of Wound	Soft tissue defect at the heel with exposed Achilles tendon*
Etiology	Avulsion at the right foot after motor-bike injury
Patient	60-year-old patient

In this case a 60-year-old patient was hospitalized with an acute avulsion after motor-bike injury. The intra-operative view revealed a defect at the heel and a 2cm exposed Achilles tendon (Fig.1). The wound was treated with 1mm MatriDerm® and unmeshed split-thickness skin graft in a One-Step Procedure. 2.5 weeks later the wound area showed a closed skin with good tendency to heal (Fig.2). The physiotherapy started at this stage. A follow up of 5 weeks p. o. demonstrated a good development with a completely stable wound also over the Achilles tendon (Fig. 3, 4).




Infected dog bite at the dorsum of the hand

Type of Wound	Infected, necrotic full-thickness wound*
Etiology	Dog bite
Patient	64-year-old female

After one month of pre-treatment of a dog bite in a general hospital (three sessions of debridement, splinting of the fascia and negative pressure wound therapy) the patient was transferred to a specialized clinic with plastic and reconstructive surgery. Pre-operative view, 4 weeks after injury: Wound bed was still necrotic (Fig. 1). Day 0: Wide and deep excision of the wound to avoid further complications (Fig.2). After preparation of an adequate wound bed, dry 1mm MatriDerm® was applied to the wound (Fig.3). After rehydration of

MatriDerm® a meshed split-thickness skin graft was applied in a One-Step Procedure (Fig.4). Fixation was performed by sutures. The wound dressing consisted of fatty gauze, bulky dressing and tight bandaging. Day 6 p. o.: First dressing change with a stable wound and excellent take of the autograft (Fig. 5). 3 months p. o.: Stable wound closure of the hand (Fig.6). 2 years follow-up: The long-term result demonstrated full range of motion of the hand and a good aesthetic outcome (Fig. 7-9).



Deglovement instep and plantar regions after motor-bike injury

Type of Wound	Deep soft tissue defect on the instep and plantar regions*
Etiology	Motor-bike injury with deglovement
Patient	45-year-old male

One week after motor-bike injury the traffic victim was transferred to a specialized trauma hospital with deep soft tissue loss on the instep and sole of the foot (Fig. 1, 2). Debridement was performed including amputation of the necrotic first toe (Fig. 3). Wound closure was achieved by using 1 mm MatriDerm® in combination with unmeshed split-thickness skin graft in a One-Step Procedure (Fig. 4, 5).

Fixation was performed with vacuum therapy for 1 week. Excellent take rate at the first dressing change (Fig. 6, 7). A small hematoma under the graft was associated with the vacuum base and can be avoided by bringing the vacuum base outside the wound area. However, this did not compromise the take. Follow-up 3 months later showed a completely healed foot with excellent elasticity of the new skin (Fig. 8).




Circular saw injury

Type of Wound	Severe soft tissue defect of the left hand*
Etiology	Circular saw injury, no co-morbidities
Patient	40-year-old male

Initial view of the circular saw injury at the hand showing a deep soft tissue defect with involvement of the auricular and the annular finger (Fig.1). Based on the bone status a partial amputation to the mid-joint of the auricular finger was necessary. The wound closure was done with a retrograde pedicular forearm flap. Three days later the flap suffered from a necrosis caused by a septal vascular anomaly (Fig.2). 13 days after injury: Second surgery was required due to the flap loss.

Day 0: Preparation and coverage of the wound with a new local flap complemented by 1 mm MatriDerm® in combination with unmeshed split-thickness skin graft in a One-Step Procedure (Fig.3,4). 5 days p. o.: First dressing change showing a little wound edge necrosis of the local flap and good take of the split-skin transplant with vital coloration (Fig.5). 4 weeks p. o.: Secondary healing of the flap and stable wound closure (Fig.6). 6 months p. o.: Good wound closure with normal skin elasticity and functionality in the reconstructed area (Fig.7, 8).





Trauma by car accident

Trauma

Type of Wound	Laceration of the left lower leg and foot*
Etiology	Trauma by car accident
Patient	55-year-old female

This 55-year-old patient suffered a laceration of the left lower leg and foot (Fig. 1). After operative debridement, there was significant soft tissue loss (Fig. 2). Defect coverage was performed with a free latissimus dorsi-serratus flap to replace the soft tissue loss. On top of the muscle flap a dry MatriDerm® (1 mm) was placed (Fig. 3) and rehydrated in situ. A meshed split-skin graft (meshed 1:1.5) was placed on top of the rehydrated MatriDerm® in a One-Step Procedure.

Fixation was performed by Negative Pressure Wound Therapy (NPWT) for 5 days. At the follow-up visit 4 weeks later, stable wound closure with complete integration of the skin graft was observed (Fig. 4). Walking training was performed beginning with 14 days p.o.. Good functional and aesthetic (Fig. 5) result was achieved. Fig. 6 and 7: 24 months after injury. The transplanted skin was pliable and soft. Both, the surgeon and the patient, were satisfied with the functional and aesthetic outcome.



Chronic venous ulcer with 3.5 years follow-up

Chronic Wounds

Type of Wound	Chronic venous ulcer*
Etiology	Venous insufficiency with 5-year open wound on the medial and lateral side of the lower left leg
Patient	51-year-old male

On first examination the wound bed appeared fibrinous and was covered with a yellowish layer, but there were no signs of necrotic tissue (Fig. 1). The wound bed was treated with non-adherent gauze for 6 months prior to surgery. Antibiotics were administered and the wound was cleansed once with chlorhexidine.

During the surgery the wound bed was properly debrided (Fig. 2) prior to dry application of 1 mm MatriDerm®. The matrix was rehydrated inside the wound bed (Fig. 3). MatriDerm® was covered in a One-Step Procedure with a 0.010 inch, unmeshed, fenestrated split-thickness skin graft (STSG) (Fig. 4).

MatriDerm® and the STSG were fixated with 6 layers of fatty gauze and V.A.C. therapy with the GranuFoam dressing was applied for 5 days (100 mmHg, continuous negative pressure) (Fig. 5).

At 14 days p. o. the take rate of the STSG was 95% and at 1 month the skin graft was completely integrated with full and stable wound closure, and the patient was able to wear compression socks for his venous insufficiency.

3.5 years follow-up showed very stable wound closure with good aesthetic outcome. The wound edges are leveled with the surrounding healthy tissue (Fig. 6) and the patient has full functional results with extension and flexion of his leg (Fig. 7, flexion). Excellent wound elasticity and pliability was achieved (Fig. 8).



* Courtesy of F. Tostes, MD, Rio Grande do Sul, Brazil

Diabetic foot syndrome with gangrene

Chronic Wounds

Type of Wound	Gangrene of the right foot*
Etiology	Diabetic foot syndrome in an obese patient
Patient	60-year-old male



A 60-year-old male with diabetic foot syndrome and a history of obesity and coronary heart disease, presented with a gangrenous right foot (Fig. 1). An emergency bedside debridement was performed and IV antibiotics were started. At the following surgical debridement, the 5th metatarsal bone was resected. The wound was further debrided with the use of maggots. The wound bed was prepared with gauze-based Negative Pressure Wound Therapy (NPWT), which was set at 120 mmHg (Fig. 2).

At the time of wound closure, the excess granulation tissue was removed and a dry sheet of 1 mm MatriDerm® was

applied on the wound bed. MatriDerm® was then moistened with Ringer's lactate (Fig. 3). A split-thickness skin graft immediately followed in a One-Step Procedure (Fig. 4).

A non-adherent wound contact layer was placed over the skin graft and the graft was fixated with NPWT. After 7 days, NPWT was ceased and the skin graft demonstrated good take rate and healing (Fig. 5). The wound was dressed with fatty gauze dressings.

Two months post-operatively, wound healing had progressed well (Fig. 6). Three and six months p. o., the wound showed stable wound closure and the patient was able to walk with a diabetic shoe (Fig. 7, 8).



Chronic diabetic ulcer at clubfoot



Chronic Wounds

Type of Wound	Chronic diabetic foot ulcer*
Etiology	Neuropathic cavoid foot; idiopathic clubfoot; chronic infected ulceration
Patient	47-year-old male

This patient with a neuropathic cavoid foot and an idiopathic clubfoot has a 4-year history of infected ulcerations on the plantar and lateral dorsal side of the foot. In the weeks prior to the reconstructive surgery, the Pseudomonas-infected ulcer was treated with MediHoney to control the infection. The wound bed was optimized with Hyiodine.

At the day of surgery, the infection was resolved and the wound size was reduced. The ulcer was thoroughly debrided resulting in a well vascularized wound bed (Fig.1,2). MatriDerm® was applied and in a One-Step Procedure covered with a meshed split-thickness skin graft which was sutured to the wound edges

(Fig.3). Negative pressure wound therapy was initiated to immobilize the graft (Fig.4). At the fifth p. o. day, signs of initial graft integration were observed (Fig.5). At 6 months p. o. the wound was completely closed with good tissue elasticity and mobility (Fig.6). The patient was pain free.



* Courtesy of H. Mössner, MD, Salzburg, Austria

Medihoney by Derma Sciences, Inc

Hyiodine by Contipro Group s.r.o.

Chronic mixed ulcer in diabetic patient

Type of Wound	Chronic mixed ulcer of the forefoot*
Etiology	Diabetic patient with mixed ulcers repeatedly covered with split skin
Patient	70-year-old patient

The 70-year-old patient suffers from a mixed chronic ulcers. The wound bed was debrided and conditioned for 2–3 weeks (Fig. 1).

Afterwards 1 mm MatriDerm® was applied in a One-Step Procedure (Fig. 2) with unmeshed split-thickness skin on top of the matrix (Fig. 3). After fixation with a vacuum system the wound was closed in 1 week after the surgical procedure (Fig. 4). Fig. 5 shows a stable result already in week 2 p. o.



Large pressure ulcer after an ischaemic stroke

Type of Wound	Large pressure ulcer of the left thorax and the lateral side of the left knee*
Etiology	Ischaemic stroke, lying on the floor for 4 days
Patient	70-year-old male

A 70-year-old patient suffered from an ischaemic stroke at home. He was found 4 days later lying on the floor unable to move and had developed large pressure ulcers of the left thorax and the lateral side of the left knee. After cardiopulmonary stabilisation at the intensive care unit, operative debridement of the necrotic tissue exposed both knee capsule and rib cartilage.

Fig. 1, 2 shows the left knee and left side of the thorax one week after initial debridement and the use of negative pressure wound therapy (NPWT) to obtain proper granulation of the wound beds.

Defect coverage was performed in a One-Step Procedure with 1 mm MatriDerm® and unmeshed split skin grafts. NPWT was then used to encourage optimum fixation of the MatriDerm® and split-thickness skin grafts. This was discontinued after one week.

Besides optimal graft take, MatriDerm® provided reliable defect coverage (Fig. 3, 4).



Large squamous cell carcinoma


Cancer Excision

Type of Wound	Squamous cell carcinoma in the left popliteal fossa*
Etiology	Growing ulceration of 5 years
Patient	81-year-old male

An 81-year-old male patient presented with a squamous cell carcinoma in the left popliteal fossa. The ulceration was growing already for 5 years.

The tumor was excised down to the crural fascia with a 1.5cm margin (Fig. 1). Negative Pressure Wound Therapy (NPWT) was initiated to prepare the wound for closure. Two weeks after the tumor excision, the wound showed a healthy granulated wound bed (Fig. 2). At the following surgical procedure, the wound was refreshed and a dry sheet of 1 mm MatriDerm® was placed on the wound bed and moistened with Ringer's lactate (Fig. 3). MatriDerm® was then trimmed to fit the wound edges. A 0.3mm split-thickness skin graft was positioned on top

of MatriDerm® in a One-Step Procedure (Fig. 4). NPWT was used for 5 days to fixate the graft. At the time of the NPWT dressing removal, the wound bed showed good signs of early healing progression (Fig. 5). The wound was dressed with a fatty gauze dressing as well as local foam compression and the knee was immobilized for 2 weeks. At the follow-up visits, the wound showed stable wound healing (Fig. 6: 1 month, Fig. 7: 3 months). At 9 months, the wound was still closed and stable, with full functional capacity (Fig. 8). The patient was highly satisfied without any functional deficit or discomfort.



Basal cell carcinoma excision in face

Cancer Excision

Type of Wound	Surgical full-thickness wound*
Etiology	Basal cell carcinoma
Patient	42-year-old female

Day 0: R0-excision of the carcinoma (Fig. 1). The intra-operative view showing the dry application of 1 mm MatriDerm® on the wound bed fixed with normal sutures (Fig. 2). Immediate coverage of MatriDerm® with a fenestrated split-thickness skin graft in a One-Step Procedure (Fig. 3). Afterwards the wound dressing was performed by using fatty gauze, bulky dressing and tight bandaging.

Day 14 p. o.: The post-operative view shows a complete wound closure with a full graft take of the transplant (Fig. 4). Follow-up 7 months: The reconstructed skin demonstrated excellent elasticity and pliability, characteristics seen by the facial movement of this skin region (Fig. 5–8). The surgeon as well as the patient were satisfied with the functional and aesthetic outcome in this special body region.



Carcinoma resection in 97-year-old patient


Cancer Excision

Type of Wound	Resection of squamous cell carcinoma*
Etiology	Fungating spinalioma cranium
Patient	97-year-old male

This 97-year-old patient, with a history of diabetes type II, presented with a fungating spinalioma on the cranium (Fig. 1). One day pre-operatively the lesion was covered with a betadine dressing.

A wide resection of the carcinoma with a 2 cm safety margin was performed. This resulted in a 6 x 7 cm defect on the frontal side of the cranium (Fig. 2). Subsequently MatriDerm® and a split-thickness skin graft were applied on the defect in a One-Step Procedure. The surgical site was dressed with vaseline gauze and foam dressing, which

were left in place for 5 days. At 1 week p. o. a 85 % take rate of the skin graft was observed, and 2 months p. o. the take rate was 100%. At the follow-up visit after one year complete healing was observed with an excellent aesthetic result (Fig. 3).




Melanoma on the nasal ridge

Cancer Excision

Type of Wound	Resection of melanoma on the nasal ridge*
Etiology	Melanoma on the nasal ridge
Patient	72-year-old female

The cancer was excised widely, resulting in a deep dermal wound down to the cartilage (Fig. 1 one week after excision). Intraoperative (Fig. 2). MatriDerm® was applied to the wound bed and trimmed appropriately. A split-thickness skin graft (STSG) was placed on top of MatriDerm® in a One-Step Procedure. The STSG was taken from the eyelids because of similar skin texture to the nasal ridge. Short-term follow-up (Fig. 3) shows viable STSG with typical pink appearance. At long-term follow-up (Fig. 4) the grafted area showed similar skin color as the surrounding skin with a very good skin elasticity (Fig. 5 and 6).



Adherent scar after extravasation of cytotoxic drugs

Adhesion Barrier


Type of Wound	Adherent scar after extravasation of cytotoxic drugs*
Etiology	Adherent scar over the flexor tendon of the hand after extravasation of cytostatics
Patient	57-year-old female

As a side-effect of chemotherapy for breast cancer treatment in 2001, extravasation led to tissue necrosis on the back of the hand. The resulting adherent scar limited hand movement and caused pain. A release of the scar tissue and a tenolysis of the extensor tendon performed in 2002 did not improve pain and mobility.

Pre-operative view: A 4 cm long scar on the back of the hand with 5 cm wide atrophic and hyperpigmented skin around it. The skin had adhered to the underlying tissue and was not movable (Fig. 1). There was a marked limitation in flexion and extension mobility (Range of Motion: Flexion 40°, extension 30°), and ulnar and radial abduction (Range of Motion: Ulnar 25°, radial 5°).

After careful detachment of the skin from the tendons, a 1 mm rehydrated sheet of MatriDerm® was inserted between the layers (Fig. 2). An intra-cutaneous continuous suture was used for closure.

Post-operatively the wound site showed good vascularization without irritation. One year p. o. the patient is completely complaint-free with good movement of the skin over the underlying tissue. There is complete mobility of the hand (Range of Motion: Flexion 80°, extension 90°, ulnar 40°, radial 38°, (Fig.3–5). The patient can clench her fist without any problems (Fig.6) and can fully extend the hand (Fig. 7). The patient is very satisfied with the functional and aesthetic result.



Treatment of adherent foot tendon

Adhesion Barrier

Type of Wound	Tenolysis*
Etiology	Tendon adhesion after foot surgery
Patient	63-year-old patient

Surgical approach at the musculus flexor digitorum brevis. Preparation and release of the adhesive tissue (Fig. 1). Gentle pull through of a small sheet of dry 1 mm MatriDerm® (Fig. 2-5). Wrapping of the tendon (Fig. 6). Final view before wound closure (Fig. 7). No reoccurrence of the claw toe and the patient no longer has an incorrect position and is able to wear his own shoes.




Adherent scar at knee

Type of Wound	Adherent scar at knee*
Etiology	Tear of cruciate ligament with adhesions
Patient	30-year-old female

A 30-year-old female patient presented with a painful scar with adhesions and limited movement after a torn cruciate ligament of the left knee. After refixation of the ligament with a titanium screw followed by several revision surgeries due to infection, adhesions had developed (Fig. 1).

In the release procedure, the scar was excised, and adhesiolysis was performed of the altered subcutaneous tissues up to the periosteum in the region of the implanted titanium screw. Two 2x4cm 2mm sheets of MatriDerm® protective matrix were placed in between the epiperiosteal and subcutaneous space, in order to prevent further adhesions. Directly on top of the rehydrated MatriDerm®, as well as in the subcutaneous tissue, fat cells were transferred that were gained with lipo-aspiration (Fig. 2).



Adhesion Barrier

The wound was sutured and covered with sterile wound dressings. Increased load bearing was initiated 2 weeks post-operatively.

One week post-operatively, the patient indicated a marked improvement in pain complaints. Three weeks post-operatively the sutures were removed and the wound showed good aesthetic result (Fig. 3). The patient reported virtually no pain and was able to fully bend the knee.

Three months post-operatively, the patient was able to put full load bearing on the knee and was free of complaints. The skin demonstrated excellent pliability (Fig. 4). This case demonstrates the successful use of MatriDerm® as a protective matrix for prevention of recurrent adhesions.



Exposed tendon of the hand

Type of Wound	Soft tissue defect with exposed tendon*
Etiology	Squamous cell carcinoma on dorsum of left hand
Patient	80-year-old female

An 80-year-old patient with a history of hypertension, was referred with a suspicious lesion on the dorsum of the left hand, which was confirmed to be a squamous cell carcinoma by histopathological examination.

The 2 x 2 cm tumor lesion was excised with appropriate margins. The wound bed was carefully debrided, which resulted in a defect with exposed tendons (Fig. 1).

The defect was covered with a 1 mm dry sheet of MatriDerm® which was rehydrated in situ (Fig. 2). MatriDerm® was then cut to fit the wound bed with a 2 mm margin (Fig. 3). A split-thickness skin graft (STSG) was then placed on top of MatriDerm® in a One-Step Procedure (Fig 4). The STSG was fixated using sutures.



Adhesion Barrier

The wound was dressed with 4 layers of non-adherent gauze, 5 layers of bulky dressing and was fixed with tight bandaging. Appropriate immobilization was ensured.

The first dressing change was performed on the 8th day after surgery and showed a stable wound bed and an excellent take rate of the graft (Fig. 5). Fifteen days post-operatively the wound was closed completely without any signs of breakdown. At 3 month follow-up the wound was healed completely with excellent pliability and full range of motion without tendon adhesions (Fig. 6–8). There was no need for further surgeries.

Donor Site after Phalloplasty with the Radial Forearm Flap

Donor Sites

Type of Wound	Donor site*
Etiology	Phalloplasty with the radial forearm free flap (RFFF) is associated with a large
Patient	female-to-male transgender patient

Consecutive patient underwent female-to-male affirming surgery. On the preoperative day, the flap was drawn on the non-dominant hand and an Allen test was performed to confirm vascular support from the ulnar artery. Under a tourniquet, the flap was raised distally to proximally and care was taken to meticulously clip all vessel branches. The radial sensory nerve (asterisk Fig. 1) was identified, dissected, and spared. The flap was raised in a subfascial plane and care was taken to include the proximal sensory nerves, veins and axial pedicle.

At this donor site, the flexor carpi radialis and brachioradialis muscle bellies were approximated and sutured together before skin graft coverage, providing a better recipient site for the graft. In addition, the flexor carpi radialis epitendineum was preserved to maximally support graft take.

After a meticulous haemostasis, two MatriDerm® sheets (1 mm) were placed over the defect (Fig. 2). An STSG was harvested using an air dermatome of 0.2mm thickness and fibrin glue¹ was sprayed on the STSG. The STSG was directly placed over collagen elastin matrix in a one-step procedure. A vacuum-assisted closure² was applied for 1 week at continuous -125mm Hg pressure (Fig. 3).

4-year after surgery the aesthetic and functional outcome demonstrate very good parameter in sensory and functional skin texture, quality and cosmetic appearance (Fig. 7). The skin could be pinched more than 1 cm (Fig. 4.)

This case shows the good sensory and functional results achieved by using MatriDerm® at the donor site after RFFF harvesting in transgender surgery.



* Courtesy of W. Watta, MD, Lausanne, Switzerland
Watta et al., MatriDerm Decreases Donor Site Morbidity After Radial Forearm Free Flap Harvest in Transgender Surgery, J Sex Med. 2017 Oct;14(10):1277-1284.

1 Artiss, Baser Healthcare Cooperation, Westlake Village, CA, USA
2 KCI, San Antonio, TX, USA

Fast Regeneration

Restored Motion


Benefits of  **matriderm**[®]
DERMAL MATRIX

- Fast integration & revascularization
- Effective closure of full-thickness injuries
- Cost-effective one-step procedure
- Superior functional & aesthetic outcomes

	Ref. No.	Size
A4	83500 - 200	210 x 297 x 1 mm
	83400 - 200	210 x 297 x 2 mm
A6	83403 - 200	105 x 148 x 1 mm
	83401 - 200	105 x 148 x 2 mm
A8	83404 - 200	52 x 74 x 1 mm

52 mm

74 mm



MedSkin Solutions Dr. Suwelack AG
Josef-Suwelack-Strasse 2
48727 Billerbeck | Germany

P +49 (0)2543 2182-0
E info@medskin-suwelack.com
W www.medskin-suwelack.com